

FACT SHEET FOR RECIPIENTS AND CAREGIVERS
EMERGENCY USE AUTHORIZATION (EUA) OF
THE MODERNA COVID-19 VACCINE TO PREVENT CORONAVIRUS DISEASE 2019
(COVID-19) IN INDIVIDUALS 18 YEARS OF AGE AND OLDER

You are being offered the Moderna COVID-19 Vaccine to prevent Coronavirus Disease 2019 (COVID-19) caused by SARS-CoV-2. This Fact Sheet contains information to help you understand the risks and benefits of the Moderna COVID-19 Vaccine, which you may receive because there is currently a pandemic of COVID-19.

The Moderna COVID-19 Vaccine is a vaccine and may prevent you from getting COVID-19. There is no U.S. Food and Drug Administration (FDA) approved vaccine to prevent COVID-19.

Read this Fact Sheet for information about the Moderna COVID-19 Vaccine. Talk to the vaccination provider if you have questions. It is your choice to receive the Moderna COVID-19 Vaccine.

The Moderna COVID-19 Vaccine is administered as a 2-dose series, 1 month apart, into the muscle.

The Moderna COVID-19 Vaccine may not protect everyone.

This Fact Sheet may have been updated. For the most recent Fact Sheet, please visit www.modernatx.com/covid19vaccine-eua.

WHAT YOU NEED TO KNOW BEFORE YOU GET THIS VACCINE

WHAT IS COVID-19?

COVID-19 is caused by a coronavirus called SARS-CoV-2. This type of coronavirus has not been seen before. You can get COVID-19 through contact with another person who has the virus. It is predominantly a respiratory illness that can affect other organs. People with COVID-19 have had a wide range of symptoms reported, ranging from mild symptoms to severe illness. Symptoms may appear 2 to 14 days after exposure to the virus. Symptoms may include: fever or chills; cough; shortness of breath; fatigue; muscle or body aches; headache; new loss of taste or smell; sore throat; congestion or runny nose; nausea or vomiting; diarrhea.

WHAT IS THE MODERNA COVID-19 VACCINE?

The Moderna COVID-19 Vaccine is an unapproved vaccine that may prevent COVID-19. There is no FDA-approved vaccine to prevent COVID-19.

The FDA has authorized the emergency use of the Moderna COVID-19 Vaccine to prevent COVID-19 in individuals 18 years of age and older under an Emergency Use Authorization (EUA).

For more information on EUA, see the “**What is an Emergency Use Authorization (EUA)?**” section at the end of this Fact Sheet.

WHAT SHOULD YOU MENTION TO YOUR VACCINATION PROVIDER BEFORE YOU GET THE MODERNA COVID-19 VACCINE?

Tell your vaccination provider about all of your medical conditions, including if you:

- have any allergies
- have a fever
- have a bleeding disorder or are on a blood thinner
- are immunocompromised or are on a medicine that affects your immune system
- are pregnant or plan to become pregnant
- are breastfeeding
- have received another COVID-19 vaccine

WHO SHOULD GET THE MODERNA COVID-19 VACCINE?

FDA has authorized the emergency use of the Moderna COVID-19 Vaccine in individuals 18 years of age and older.

WHO SHOULD NOT GET THE MODERNA COVID-19 VACCINE?

You should not get the Moderna COVID-19 Vaccine if you:

- had a severe allergic reaction after a previous dose of this vaccine
- had a severe allergic reaction to any ingredient of this vaccine

WHAT ARE THE INGREDIENTS IN THE MODERNA COVID-19 VACCINE?

The Moderna COVID-19 Vaccine contains the following ingredients: messenger ribonucleic acid (mRNA), lipids (SM-102, polyethylene glycol [PEG] 2000 dimyristoyl glycerol [DMG], cholesterol, and 1,2-distearoyl-sn-glycero-3-phosphocholine [DSPC]), tromethamine, tromethamine hydrochloride, acetic acid, sodium acetate, and sucrose.

HOW IS THE MODERNA COVID-19 VACCINE GIVEN?

The Moderna COVID-19 Vaccine will be given to you as an injection into the muscle.

The Moderna COVID-19 Vaccine vaccination series is 2 doses given 1 month apart.

If you receive one dose of the Moderna COVID-19 Vaccine, you should receive a second dose of the same vaccine 1 month later to complete the vaccination series.

HAS THE MODERNA COVID-19 VACCINE BEEN USED BEFORE?

The Moderna COVID-19 Vaccine is an unapproved vaccine. In clinical trials, approximately 15,400 individuals 18 years of age and older have received at least 1 dose of the Moderna COVID-19 Vaccine.

WHAT ARE THE BENEFITS OF THE MODERNA COVID-19 VACCINE?

In an ongoing clinical trial, the Moderna COVID-19 Vaccine has been shown to prevent COVID-19 following 2 doses given 1 month apart. The duration of protection against COVID-19 is currently unknown.

WHAT ARE THE RISKS OF THE MODERNA COVID-19 VACCINE?

Side effects that have been reported with the Moderna COVID-19 Vaccine include:

- Injection site reactions: pain, tenderness and swelling of the lymph nodes in the same arm of the injection, swelling (hardness), and redness
- General side effects: fatigue, headache, muscle pain, joint pain, chills, nausea and vomiting, and fever

There is a remote chance that the Moderna COVID-19 Vaccine could cause a severe allergic reaction. A severe allergic reaction would usually occur within a few minutes to one hour after getting a dose of the Moderna COVID-19 Vaccine. For this reason, your vaccination provider may ask you to stay at the place where you received your vaccine for monitoring after vaccination. Signs of a severe allergic reaction can include:

- Difficulty breathing
- Swelling of your face and throat
- A fast heartbeat
- A bad rash all over your body
- Dizziness and weakness

These may not be all the possible side effects of the Moderna COVID-19 Vaccine. Serious and unexpected side effects may occur. The Moderna COVID-19 Vaccine is still being studied in clinical trials.

WHAT SHOULD I DO ABOUT SIDE EFFECTS?

If you experience a severe allergic reaction, call 9-1-1, or go to the nearest hospital.

Call the vaccination provider or your healthcare provider if you have any side effects that bother you or do not go away.

Report vaccine side effects to **FDA/CDC Vaccine Adverse Event Reporting System (VAERS)**. The VAERS toll-free number is 1-800-822-7967 or report online to <https://vaers.hhs.gov/reportevent.html>. Please include “Moderna COVID-19 Vaccine EUA” in the first line of box #18 of the report form.

In addition, you can report side effects to ModernaTX, Inc. at 1-866-MODERNA (1-866-663-3762).

You may also be given an option to enroll in **v-safe**. **V-safe** is a new voluntary smartphone-based tool that uses text messaging and web surveys to check in with people who have been vaccinated to identify potential side effects after COVID-19 vaccination. **V-safe** asks questions that help CDC monitor the safety of COVID-19 vaccines. **V-safe** also provides second-dose reminders if needed and live telephone follow-up by CDC if participants report a significant health impact following COVID-19 vaccination. For more information on how to sign up, visit: www.cdc.gov/vsafe.

WHAT IF I DECIDE NOT TO GET THE MODERNA COVID-19 VACCINE?

It is your choice to receive or not receive the Moderna COVID-19 Vaccine. Should you decide not to receive it, it will not change your standard medical care.

ARE OTHER CHOICES AVAILABLE FOR PREVENTING COVID-19 BESIDES MODERNA COVID-19 VACCINE?

Currently, there is no FDA-approved alternative vaccine available for prevention of COVID-19. Other vaccines to prevent COVID-19 may be available under Emergency Use Authorization.

CAN I RECEIVE THE MODERNA COVID-19 VACCINE WITH OTHER VACCINES?

There is no information on the use of the Moderna COVID-19 Vaccine with other vaccines.

WHAT IF I AM PREGNANT OR BREASTFEEDING?

If you are pregnant or breastfeeding, discuss your options with your healthcare provider.

WILL THE MODERNA COVID-19 VACCINE GIVE ME COVID-19?

No. The Moderna COVID-19 Vaccine does not contain SARS-CoV-2 and cannot give you COVID-19.


KEEP YOUR VACCINATION CARD

When you receive your first dose, you will get a vaccination card to show you when to return for your second dose of the Moderna COVID-19 Vaccine. Remember to bring your card when you return.

ADDITIONAL INFORMATION

If you have questions, visit the website or call the telephone number provided below.

To access the most recent Fact Sheets, please scan the QR code provided below.

Moderna COVID-19 Vaccine website	Telephone number
www.modernatx.com/covid19vaccine-eua 	1-866-MODERNA (1-866-663-3762)

HOW CAN I LEARN MORE?

- Ask the vaccination provider
- Visit CDC at <https://www.cdc.gov/coronavirus/2019-ncov/index.html>
- Visit FDA at <https://www.fda.gov/emergency-preparedness-and-response/mcm-legal-regulatory-and-policy-framework/emergency-use-authorization>
- Contact your state or local public health department

WHERE WILL MY VACCINATION INFORMATION BE RECORDED?

The vaccination provider may include your vaccination information in your state/local jurisdiction's Immunization Information System (IIS) or other designated system. This will ensure that you receive the same vaccine when you return for the second dose. For more information about IISs, visit: <https://www.cdc.gov/vaccines/programs/iis/about.html>.

WHAT IS THE COUNTERMEASURES INJURY COMPENSATION PROGRAM?

The Countermeasures Injury Compensation Program (CICP) is a federal program that may help pay for costs of medical care and other specific expenses of certain people who have been seriously injured by certain medicines or vaccines, including this vaccine. Generally, a claim must be submitted to the CICP within one (1) year from the date of receiving the vaccine. To learn more about this program, visit www.hrsa.gov/cicp/ or call 1-855-266-2427.

WHAT IS AN EMERGENCY USE AUTHORIZATION (EUA)?

The United States FDA has made the Moderna COVID-19 Vaccine available under an emergency access mechanism called an EUA. The EUA is supported by a Secretary of Health and Human Services (HHS) declaration that circumstances exist to justify the emergency use of drugs and biological products during the COVID-19 pandemic.


The Moderna COVID-19 Vaccine has not undergone the same type of review as an FDA-approved or cleared product. FDA may issue an EUA when certain criteria are met, which includes that there are no adequate, approved, and available alternatives. In addition, the FDA decision is based on the totality of the scientific evidence available showing that the product may be effective to prevent COVID-19 during the COVID-19 pandemic and that the known and potential benefits of the product outweigh the known and potential risks of the product. All of these criteria must be met to allow for the product to be used during the COVID-19 pandemic.

The EUA for the Moderna COVID-19 Vaccine is in effect for the duration of the COVID-19 EUA declaration justifying emergency use of these products, unless terminated or revoked (after which the products may no longer be used).

©2020 ModernaTX, Inc. All rights reserved.

Patent(s): www.modernatx.com/patents

Revised: 12/2020



Pre-Vaccination Checklist for COVID-19 Vaccines



For vaccine recipients:

The following questions will help us determine if there is any reason you should not get the COVID-19 vaccine today.

If you answer "yes" to any question, it does not necessarily mean you should not be vaccinated. It just means additional questions may be asked. If a question is not clear, please ask your healthcare provider to explain it.

Patient Name _____

Age _____

	Yes	No	Don't know
1. Are you feeling sick today?			
2. Have you ever received a dose of COVID-19 vaccine?			
<ul style="list-style-type: none"> • If yes, which vaccine product? <ul style="list-style-type: none"> <input type="checkbox"/> Pfizer <input type="checkbox"/> Moderna <input type="checkbox"/> Another product _____ 			
3. Have you ever had a severe allergic reaction (e.g., anaphylaxis) to something? For example, a reaction for which you were treated with epinephrine or EpiPen®, or for which you had to go to the hospital?			
<ul style="list-style-type: none"> • Was the severe allergic reaction after receiving a COVID-19 vaccine? • Was the severe allergic reaction after receiving another vaccine or another injectable medication? 			
4. Have you received passive antibody therapy (monoclonal antibodies or convalescent serum) as treatment for COVID-19?			
5. Have you received another vaccine in the last 14 days?			
6. Have you had a positive test for COVID-19 or has a doctor ever told you that you had COVID-19?			
7. Do you have a weakened immune system caused by something such as HIV infection or cancer or do you take immunosuppressive drugs or therapies?			
8. Do you have a bleeding disorder or are you taking a blood thinner?			
9. Are you pregnant or breastfeeding?			

Form reviewed by _____

Date _____

This page has been left blank.

Pre-Vaccination Checklist for COVID-19 Vaccines

Information for Healthcare Professionals



For additional information on COVID-19 vaccine clinical guidance, see: <https://www.cdc.gov/vaccines/covid-19/info-by-product/clinical-considerations.html>.

For additional information on ACIP general recommendations, see: <https://www.cdc.gov/vaccines/hcp/acip-recs/general-recs/index.html>.

Two COVID-19 vaccines are currently authorized for use in the United States. These vaccines are authorized for use among different age populations.

PRODUCT	AUTHORIZED AGE GROUPS
Pfizer-BioNTech COVID-19 Vaccine	16 years of age and older
Moderna COVID-19 Vaccine	18 years of age and older

Anyone outside of the authorized age groups for a product should not receive the vaccine.

Are you feeling sick today?

There is no evidence that acute illness reduces vaccine efficacy or increases vaccine adverse events. However, as a precaution with moderate or severe acute illness, all vaccines should be delayed until the illness has improved. **Mild illnesses (e.g., upper respiratory infections, diarrhea) are NOT contraindications to vaccination.** Do not withhold vaccination if a person is taking antibiotics.

Vaccination of persons with current SARS-CoV-2 infection should be deferred until the person has recovered from acute illness and they can discontinue isolation. This recommendation applies to persons who develop SARS-CoV-2 infection before receiving any vaccine doses as well as those who develop SARS-CoV-2 infection after the first dose but before receipt of the second dose.

Have you ever received a dose of COVID-19 vaccine?

COVID-19 vaccines are **NOT** interchangeable. Currently authorized COVID-19 vaccines require two doses. Both doses of the series should be completed with the same product. Product dosing schedules vary.

Check medical records, immunization information systems, and vaccination record cards to help determine the initial product received. Those who received a trial vaccine should consult with the trial sponsors to determine if it is feasible to receive additional doses.

PRODUCT	DOSING SCHEDULE Between doses 1 and 2
Pfizer-BioNTech COVID-19 Vaccine	21 days
Moderna COVID-19 Vaccine	28 days

The second dose should be administered as close to the recommended interval as possible. The vaccine can be given up to four days in advance of the recommended interval if a patient presents early and you are concerned they will not return at the appropriate interval for vaccination. However, there is no maximum interval between the first and second dose for either vaccine. The series does not need to be restarted.

Pre-Vaccination Checklist for COVID-19 Vaccines

Information for Healthcare Professionals



COVID-19 Vaccine Components

Description	Pfizer-BioNTech COVID-19 vaccine	Moderna COVID-19 vaccine
mRNA	Nucleoside-modified mRNA encoding the viral spike (S) glycoprotein of SARS-CoV-2	Nucleoside-modified mRNA encoding the viral spike (S) glycoprotein of SARS-CoV-2
Lipids	2[(polyethylene glycol)-2000]-N, N-ditetradecylacetamide	Polyethylene glycol (PEG) 2000 dimyristoyl glycerol (DMG)
	1,2-distearoyl-sn-glycero-3-phosphocholine	1,2-distearoyl-sn-glycero-3-phosphocholine
	Cholesterol	Cholesterol
	(4-hydroxybutyl)azanediyl)bis(hexane-6,1-diyl) bis(2-hexyldecanoate)	SM-102 (Proprietary to Moderna)
Salts, sugars, buffers	Potassium chloride	Tromethamine
	Monobasic potassium phosphate	Tromethamine hydrochloride
	Sodium chloride	Acetic acid
	Dibasic sodium phosphate dihydrate	Sodium acetate
	Sucrose	Sucrose

Have you ever had a severe allergic reaction (e.g., anaphylaxis) to something? For example, a reaction for which you were treated with epinephrine or EpiPen®, or for which you had to go to the hospital?

Allergic reactions, including severe allergic reactions, NOT related to vaccines or injectable therapies (e.g., food, pet, venom, environmental, or latex allergies; oral medications) are NOT a contraindication or precaution to vaccination with currently authorized COVID-19 vaccine. HOWEVER, individuals who have had severe allergic reactions to something, regardless of cause, **should be observed for 30 minutes after vaccination**. All other persons should be observed for 15 minutes.

Was the severe allergic reaction after receiving a COVID-19 vaccine?

History of severe allergic reaction (e.g., anaphylaxis) to a previous dose or component of the COVID-19 vaccine product being offered is a contraindication to any current COVID-19 vaccine. Ask questions about previous severe reactions that might indicate an allergy to a vaccine component. For example, PEG may have been a component of medication for a colonoscopy.

Was the severe allergic reaction after receiving another vaccine or another injectable medication?

History of severe allergic reaction (e.g., anaphylaxis) to another vaccine or a component of another vaccine OR anaphylactic reaction to any other injectable medication is a **precaution to currently authorized COVID-19 vaccine**. Vaccine may be given, but counsel patients about unknown risks of developing a severe allergic reaction and balance these risks against the benefits of vaccination. These individuals should be observed for 30 minutes after vaccination. A history of mild allergic reaction to a vaccine or injectable therapy is not a precaution to vaccination.

Healthcare professionals should be familiar with identifying immediate-type allergic reactions, including anaphylaxis, and be competent in treating these events at the time of vaccine administration. Appropriate medical treatment for severe allergic reactions must be immediately available in the event that an acute anaphylactic reaction occurs following administration of a COVID-19 vaccine.

See [Management of Anaphylaxis at COVID-19 Vaccination Sites](#) | CDC for additional guidance.

Have you received passive antibody therapy as treatment for COVID-19?

Based on the estimated half-life of monoclonal antibodies or convalescent plasma as part of COVID-19 treatment, as well as evidence suggesting that reinfection is uncommon in the 90 days after initial infection, **vaccination should be deferred for at least 90 days**, as a precautionary measure until additional information becomes available, to avoid interference of the antibody treatment with vaccine-induced immune responses.

Pre-Vaccination Checklist for COVID-19 Vaccines

Information for Healthcare Professionals



Clinical Consideration Questions

Responses to these questions are not (on their own) contraindications or precautions to vaccination. However, healthcare professionals should be prepared to discuss information and options with patients based on their responses to the following questions.

Have you received another vaccine in the last 14 days?

COVID-19 vaccine series should be administered alone, with a minimum interval of 14 days before or after administration with other vaccines. This recommendation is based on the lack of data on the safety and efficacy of mRNA COVID-19 vaccines administered simultaneously with other vaccines.

Have you had a positive test for COVID-19 or has a doctor ever told you that you had COVID-19?

Vaccination should be offered to persons regardless of history of prior symptomatic or asymptomatic SARS-CoV-2 infection. Vaccination of persons with known current SARS-CoV-2 infection should be deferred until the person has recovered from the acute illness (if the person had symptoms) and criteria have been met for them to discontinue isolation.

Persons with documented acute SARS-CoV-2 infection in the preceding 90 days may delay vaccination until near the end of this period, if desired, because current evidence suggests reinfection is uncommon during this time.

Viral testing to assess for acute SARS-CoV-2 infection or serologic testing to assess for prior infection solely for the purposes of vaccine decision-making is not recommended.

Do you have a weakened immune system caused by something such as HIV infection or cancer or do you take immunosuppressive drugs or therapies?

Persons with HIV infection or other immunocompromising conditions, or who take immunosuppressive medications or therapies might be at increased risk for severe COVID-19. mRNA COVID-19 vaccines may be administered to persons with underlying medical conditions who have no contraindications to vaccination. However, they should be counseled about the unknown vaccine safety profile and effectiveness in immunocompromised populations, as well as the potential for reduced immune responses and the need to continue to follow all current guidance to protect themselves against COVID-19, including wearing a mask, social distancing, and washing hands frequently.

Do you have a bleeding disorder or are you taking a blood thinner?

COVID-19 vaccine may be given to these patients, if a physician familiar with the patient's bleeding risk determines that the vaccine can be administered intramuscularly with reasonable safety. ACIP recommends the following technique for intramuscular vaccination in patients with bleeding disorders or taking blood thinners: a fine-gauge needle (23-gauge or smaller caliber) should be used for the vaccination, followed by firm pressure on the site, without rubbing, for at least 2 minutes.

Are you pregnant or breastfeeding?

If pregnant people are part of a group that is recommended to receive a COVID-19 vaccine (e.g., healthcare personnel), they may choose to be vaccinated. For pregnant people seeking guidance in making a decision, pregnant people and their healthcare providers should consider the level of COVID-19 community transmission, the patient's personal risk of contracting COVID-19, the risks of COVID-19 to the patient and potential risks to the fetus, the efficacy of the vaccine, the side effects of the vaccine, and the lack of data about the vaccine during pregnancy.

A lactating person who is part of a group recommended to receive a COVID-19 vaccine (e.g., healthcare personnel) may choose to be vaccinated. There are no data on the safety of COVID-19 vaccines in lactating people or the effects of mRNA COVID-19 vaccines on the breastfed infant or milk production/excretion.



ImmTrac2 Immunization Registry
DISASTER INFORMATION
RETENTION CONSENT FORM



(Please print clearly)

Client's Last Name grid

Client's Last Name

Client's First Name grid

Client's First Name

Client's Middle Name grid

Client's Middle Name

Client's Date of Birth grid

Client's Date of Birth

*A parent, legal guardian or managing conservator must sign this form if the client is younger than 18 years of age.

Client's Gender: Male Female

Client's Address grid

Client's Address

Apartment # grid

Apartment #

Client's Telephone grid

Client's Telephone

City grid

City

State grid

State

Zip Code grid

Zip Code

County grid

County

Mother's First Name grid

Mother's First Name (if client is younger than 18 years of age)

Mother's Maiden Name grid

Mother's Maiden Name (if client is younger than 18 years of age)

ImmTrac2, the Texas immunization registry, has been designated as the disaster-related reporting and tracking system for immunizations, antivirals, and other medications administered to individuals in preparation for, or in response to, a disaster or public health emergency. From the time the event is declared over, ImmTrac2 will retain disaster-related information received from health-care providers for a period of 5 years. At the end of the 5 year retention period, client-specific disaster-related information will be removed from the Registry unless consent is granted to retain the client information in ImmTrac2 beyond the 5 year retention period.

The Texas Department of State Health Services (DSHS) encourages your voluntary participation in the Texas immunization registry.

Consent for Retention of Disaster-Related Information and Release of Information to Authorized Entities

I understand that, by granting the consent below, I am authorizing retention of my (or my child's) disaster-related information by DSHS beyond the 5 year retention period. I further understand that DSHS will include this information in the state's central immunization registry ("ImmTrac2"). Once in ImmTrac2, my (or my child's) disaster-related information may by law be accessed by:

- a state agency, for the purpose of aiding and coordinating communicable disease prevention and control efforts, and / or
a physician or other health-care provider legally authorized to administer immunizations, antivirals, and other medications, for treating the client as a patient;

I understand that I may withdraw this consent to retain information in the ImmTrac2 Registry beyond the 5 year retention period and my consent to release information from the Registry, at any time by written communication to the Texas Department of State Health Services, ImmTrac2 Group - MC 1946, P. O. Box 149347, Austin, Texas 78714-9347.

By my signature below, I GRANT consent to retain my disaster-related information (or my child's information if younger than age 18) in the Texas immunization registry beyond the 5 year retention period.

Client (or parent, legal guardian, or managing conservator): Printed Name:

Date: Signature:

Privacy Notification: With few exceptions, you have the right to request and be informed about information that the State of Texas collects about you. You are entitled to receive and review the information upon request. You also have the right to ask the state agency to correct any information that is determined to be incorrect. See http://www.dshs.texas.gov for more information on Privacy Notification. (Reference: Government Code, Section 552.021, 552.023, 559.003, and 559.004)

Upon completion, please fax or mail form to the DSHS ImmTrac2 Group or a registered Health-care provider.

Questions? (800) 252-9152 • (512) 776-7284 • Fax: (866) 624-0180 • www.ImmTrac.com • ImmTrac2 DC Texas Department of State Health Services • ImmTrac2 Group - MC 1946 • P. O. Box 149347 • Austin, TX 78714-9347

PROVIDERS REGISTERED WITH ImmTrac2

Please enter client information in ImmTrac2 and affirm that consent has been granted.

DO NOT fax to ImmTrac2. Retain this form in your client's record.

ImmTrac2 Immunization Registry
FORMULARIO DE CONSENTIMIENTO
DE RETENCIÓN DE INFORMACIÓN
SOBRE EL DESASTRE

(Favor de escribir claramente con letra de molde)

Apellido del Cliente**Nombre del Cliente****Fecha de Nacimiento del Cliente**

*Uno de los padres, el tutor legal o el custodio administrador debe firmar este formulario si el cliente es menor de 18 años de edad.

Dirección del Cliente**Ciudad****Nombre de la Madre (si el cliente es menor de 18 años de edad)****Segundo Nombre del Cliente****Apartamento #****Estado****Teléfono del cliente****Condado****Apellido de Soltera de la Madre (si el cliente es menor de 18 años de edad)**

Nombre de la Madre (si el cliente es menor de 18 años de edad) Apellido de Soltera de la Madre (si el cliente es menor de 18 años de edad) ImmTrac2, el registro de inmunización de Texas, ha sido designado como el sistema de información y seguimiento para inmunizaciones, antivirales y otros medicamentos administrados a individuos durante la preparación o respuesta a un desastre o emergencia de salud pública. A partir del momento en que se declare finalizado el evento, ImmTrac2 retendrá la información relacionada con el desastre recibida por profesionales de la salud durante un periodo de 5 años. Al final del periodo de retención de 5 años, la información del cliente relacionada con el desastre se removerá del registro a menos que se dé el consentimiento para retener la información en ImmTrac2 más allá del periodo de retención de 5 años.

El Departamento Estatal de Servicios de Salud de Texas (DSHS) le anima a participar voluntariamente en el registro de inmunización de Texas.

Consentimiento de retención de información relacionada con el desastre y dar a conocer la información a entidades autorizadas

Entiendo que, al dar mi consentimiento a continuación, autorizo al DSHS a que retenga mi información relacionada con el desastre (o la de mi hijo[a]) más allá del periodo de retención de 5 años. Además entiendo que el DSHS incluirá esta información en el registro central de inmunización del estado ("ImmTrac2"). Una vez que mi información relacionada con el desastre (o la de mi hijo[a]) esté en ImmTrac2, puede ésta por ley ser accedida por:

- las agencias estatales, con el propósito de ayudar con los esfuerzos de prevención y control de enfermedades transmisibles y su coordinación y
- los médicos o demás profesionales de la salud legalmente autorizados para administrar vacunas, antivirales y otros medicamentos, para el tratamiento del cliente como paciente.

Entiendo que puedo retirar el consentimiento para retener información en el registro ImmTrac2 más allá del periodo de retención de 5 años y mi consentimiento para dar a conocer la información del registro, en cualquier momento mediante comunicación escrita dirigida a Texas Department of State Health Services, ImmTrac2 Group – MC 1946, P. O. Box 149347, Austin, Texas 78714-9347.

Con mi firma a continuación, DOY mi consentimiento para que se retenga mi información relacionada con el desastre (o la de mi hijo[a] si es menor de 18 años de edad) en el registro de inmunizaciones de Texas más allá del periodo de retención de 5 años.

Cliente (o padre / madre, tutor legal o custodio administrador): _____
Nombre en letra de molde


Fecha: _____ Firma: _____

Notificación Sobre Privacidad: Tan solo por unas cuantas excepciones, usted tiene el derecho de solicitar y de ser informado sobre la información que el Estado de Texas reúne sobre usted. A usted se le debe conceder el derecho de recibir y revisar la información al requerirla. Usted también tiene el derecho de pedir que la agencia estatal corrija cualquier información que se ha determinado sea incorrecta. Diríjase a <http://www.dshs.texas.gov> para más información sobre la Notificación sobre privacidad. (Referencia: Government Code, sección 552.021, 552.023, 559.003 y 559.004)

Al llenarlo, mándelo por fax o correo postal al Grupo ImmTrac2 del DSHS o a un proveedor de salud inscrito.

¿Tiene preguntas? (800) 252-9152 • (512) 776-7284 • Fax: (866) 624-0180 • www.ImmTrac.com • ImmTrac2 DC
Texas Department of State Health Services • ImmTrac2 Group – MC 1946 • P. O. Box 149347 • Austin, TX 78714-9347

PROVIDERS REGISTERED WITH ImmTrac2Please enter client information in ImmTrac2 and **affirm** that consent has been granted.**DO NOT** fax to ImmTrac2. **Retain this form in your client's record.**



**Get vaccinated.
Get your smartphone.
Get started with v-safe.**

What is v-safe?

V-safe is a smartphone-based tool that uses text messaging and web surveys to provide personalized health check-ins after you receive a COVID-19 vaccination. Through **v-safe**, you can quickly tell CDC if you have any side effects after getting the COVID-19 vaccine. Depending on your answers, someone from CDC may call to check on you. And **v-safe** will remind you to get your second COVID-19 vaccine dose if you need one.

Your participation in CDC's **v-safe** makes a difference—it helps keep COVID-19 vaccines safe.

How can I participate?

Once you get a COVID-19 vaccine, you can enroll in **v-safe** using your smartphone. Participation is voluntary and you can opt out at any time. You will receive text messages from **v-safe** around 2pm local time. To opt out, simply text "STOP" when **v-safe** sends you a text message. You can also start **v-safe** again by texting "START."

How long do v-safe check-ins last?

During the first week after you get your vaccine, **v-safe** will send you a text message each day to ask how you are doing. Then you will get check-in messages once a week for up to 5 weeks. The questions **v-safe** asks should take less than 5 minutes to answer. If you need a second dose of vaccine, **v-safe** will provide a new 6-week check-in process so you can share your second-dose vaccine experience as well. You'll also receive check-ins 3, 6, and 12 months after your final dose of vaccine.

Is my health information safe?

Yes. Your personal information in **v-safe** is protected so that it stays confidential and private.*

*To the extent **v-safe** uses existing information systems managed by CDC, FDA, and other federal agencies, the systems employ strict security measures appropriate for the data's level of sensitivity. These measures comply, where applicable, with the following federal laws, including the Privacy Act of 1974; standards enacted that are consistent with the Health Insurance Portability and Accountability Act of 1996 (HIPAA); the Federal Information Security Management Act, and the Freedom of Information Act.



Use your smartphone to tell CDC about any side effects after getting the COVID-19 vaccine. You'll also get reminders if you need a second vaccine dose.



Sign up with your smartphone's browser at vsafe.cdc.gov

OR

Aim your smartphone's camera at this code




How to register and use v-safe

You will need your smartphone and information about the COVID-19 vaccine you received. This information can be found on your vaccination record card; if you cannot find your card, please contact your healthcare provider.

Register

1. Go to the **v-safe** website using one of the two options below:



2. Read the instructions. Click **Get Started**.
3. Enter your name, mobile number, and other requested information. Click **Register**.
4. You will receive a text message with a verification code on your smartphone. Enter the code in **v-safe** and click **Verify**.
5. At the top of the screen, click **Enter your COVID-19 vaccine information**.
6. Select which COVID-19 vaccine you received (found on your vaccination record card; if you cannot find your card, please contact your healthcare provider). Then enter the date you were vaccinated. Click **Next**.
7. Review your vaccine information. If correct, click **Submit**. If not, click **Go Back**.
8. **Congrats! You're all set!** If you complete your registration before 2pm local time, **v-safe** will start your initial health check-in around 2pm that day. If you register after 2pm, **v-safe** will start your initial health check-in immediately after you register—just follow the instructions.

You will receive a reminder text message from **v-safe** when it's time for the next check-in—around 2pm local time. Just click the link in the text message to start the check-in.

Complete a v-safe health check-in

1. When you receive a **v-safe** check-in text message on your smartphone, click the link when ready.
2. Follow the instructions to complete the check-in.

Troubleshooting

How can I come back and finish a check-in later if I'm interrupted?

- Click the link in the text message reminder to restart and complete your check-in.

How do I update my vaccine information after my second COVID-19 vaccine dose?

- **V-safe** will automatically ask you to update your second dose information. Just follow the instructions.

Need help with v-safe?

Call 800-CDC-INFO (800-232-4636)

TTY 888-232-6348

Open 24 hours, 7 days a week

Visit www.cdc.gov/vsafe

